

(19) World Intellectual Property
Organization
International Bureau



(43) International Publication Date
19 May 2005 (19.05.2005)

PCT

(10) International Publication Number
WO 2005/044775 A1

(51) International Patent Classification⁷: **C07C 67/08**,
69/017, 69/16, C07D 311/72

(21) International Application Number:
PCT/EP2004/012058

(22) International Filing Date: 26 October 2004 (26.10.2004)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
03025513.7 7 November 2003 (07.11.2003) EP

(71) Applicant (for all designated States except US): **DSM
IP ASSETS B.V.** [NL/NL]; Het Overloon 1, NL-6411 Te
Heerlen (NL).

(72) Inventors; and

(75) Inventors/Applicants (for US only): **AQUINO, Fabrice**
[FR/FR]; 34a Rue du Leimbach, F-68950 Reiningue (FR).
BONRATH, Werner [DE/DE]; 29 Luckenbachweg,
79115 Freiburg (DE). **PACE, Francesco** [IT/CH]; Salzbo-
denstrasse 6, CH-4310 Rheinfelden (CH).

(74) Agents: **GROSSNER, Lutz** et al.; c/o DSM Nutritional
Products Ltd., Wurmisweg 576, CH-4303 Kaiseraugst
(CH).

(81) Designated States (unless otherwise indicated, for every
kind of national protection available): AE, AG, AL, AM,
AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN,
CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI,
GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE,
KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD,
MG, MK, MN, MW, MX, MZ, NA, NI, NO, NZ, OM, PG,
PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SY, TJ, TM,
TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM,
ZW.

(84) Designated States (unless otherwise indicated, for every
kind of regional protection available): ARIPO (BW, GH,
GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM,
ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM),
European (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI,
FR, GB, GR, HU, IE, IT, LU, MC, NL, PL, PT, RO, SE, SI,
SK, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ,
GW, ML, MR, NE, SN, TD, TG).

Published:

— with international search report

For two-letter codes and other abbreviations, refer to the "Guid-
ance Notes on Codes and Abbreviations" appearing at the begin-
ning of each regular issue of the PCT Gazette.

(54) Title: PROCESS FOR THE PREPARATION OF 2,3,5-TRIMETHYLHYDROQUINONE DIACYLATES

(57) Abstract: 2,3,5-Trimethyl-1,4-hydroquinone diacylates are obtained by reacting 3,5,5-trimethyl-1,4-benzoquinone with an acy-
lating agent in the presence of methane trisulfonic acid as a catalyst.



WO 2005/044775 A1

Process for the preparation of 2,3,5-trimethylhydroquinone diacylates

The present invention is concerned with a process for the preparation of 2,3,5-trimethylhydroquinone diacylates by reacting 3,5,5-trimethyl-1,4-benzoquinone (ketoisophorone) with an acylating agent in the presence of methanetrissulfonic acid. 2,3,5-Trimethylhydroquinone diacylates are useful as reactants for the preparation of 2,3,5-trimethylhydroquinone, itself a known valuable reactant for the preparation of (all-rac)- α -tocopherol.

2,3,5-Trimethylhydroquinone diacylates are known to be producible by reacting ketoisophorone with an acylating agent in the presence of a strongly acidic catalyst. Many such catalysts have been proposed in the past for this purpose, in particular protonic acids, e.g. such inorganic acids as sulphuric acid; such organic acids as p-toluenesulphonic acid; strongly acidic ion exchange resins; and such Lewis acids as zinc chloride, boron trifluoride, antimony pentafluoride and titanium tetrachloride: see inter alia German Offenlegungsschrift 2149159 and European Patent Publications EP 0916642 A1 and EP 1028103 A1; as well as NH-acidic or CH-acidic catalysts, see PCT Publication WO 03/051812.

It has now been found that by the use of small amounts of methanetrissulfonic acid, the conversion of ketoisophorone to 2,3,5-trimethylhydroquinone diacylates can be accomplished in high yield and without the need to use additional solvents. The use of methanetrissulfonic acid as the catalyst provides, all in all, advantages over catalysts used so far in this reaction in terms of stability of the catalyst, high yield, selectivity, and costs.

Thus, the present invention relates to a process for the preparation of 2,3,5-trimethyl-1,4-hydroquinone diacylates by reacting 3,5,5-trimethyl-1,4-benzoquinone with an acylating agent in the presence of methane trissulfonic acid.

The acylating agent used in the process of the present invention may be any acylating agent that is conventionally used in the conversion of ketoisophorone to 2,3,5-trimethylhydroquinone acylates, particularly acid anhydrides, acyl halides, and enol esters. Examples of acid anhydrides are straight or branched chain alkanolic acid anhydrides such as acetic, propionic and butyric anhydride. Examples of acyl halides are straight or branched chain alkanoyl chlorides such as acetyl, propionyl and butyryl chloride. Finally, examples of enol esters are isopropenyl acetate and butyrate. The preferred acylating agent is acetic anhydride or acetyl chloride, especially acetic anhydride.

The process of the present invention can be carried out in the absence of a solvent. While the ratio of acylating agent to ketoisophorone is not narrowly critical the molar ratio of acylating agent to ketoisophorone is suitably from about 1 : 1 to about 10 : 1, preferably from about 5 : 1 to about 3 : 1, and is most preferably about 3 : 1.

The amount of catalyst, methane trisulfonic acid, is suitably about 0.01 to about 2.0 mole%, preferably about 0.075 to about 1.5 mole%, and most preferably about 0.1 to about 1.0 mole%, based on the amount of ketoisophorone.

The process is conveniently carried out at temperatures from about 0 °C to about 140 °C, preferably from about 20 °C to about 90 °C, especially 20 °C to 70 °C.

The process according to the present invention may be carried out batchwise or in continuous mode. Moreover, the process is conveniently carried out under an inert gas atmosphere, preferably under gaseous nitrogen or argon.

The progress of the reaction is suitably monitored by gas chromatography and mass spectrometry of samples taken from the reaction mixture at various time intervals during the reaction.

The produced 2,3,5-trimethylhydroquinone diacylate can be isolated after distilling off the remaining acylating agent and the secondary product formed in the acylation, e.g. acetic acid when acetic anhydride is used as the acylating agent, by extraction of the crude product mixture with a suitable organic solvent, e.g. toluene. For instance, in effecting this procedure using acetic anhydride as the acylating agent 2,3,5-trimethylhydroquinone diacetate was obtained as colourless crystals after evaporating off the toluene used as the extracting solvent. Another isolation procedure is the crystallization of the 2,3,5-trimethylhydroquinone diacylate from the mixture at the termination of the reaction by cooling, and, optionally, adding water, to the mixture to promote the crystallization.

The catalyst can be recovered by extraction with water or acid-water and concentration of the extract. Alternatively, the catalyst can be recovered by adding a

biphasic solvent system, e. g. a carbonate (particularly ethylene carbonate or propylene carbonate) and an aliphatic hydrocarbon (particularly heptane or octane), and isolating it from the polar (carbonate) phase

The 2,3,5-trimethylhydroquinone diacylate obtained by the process of the present invention can be converted into 2,3,5-trimethylhydroquinone by transesterification, i.e. by treatment with an alcohol, e.g. an aliphatic alcohol such as isopropanol or n-butanol. Depending on the amounts of alcohol and catalyst and on the temperature in the reaction mixture, the transesterification yields the unesterified 2,3,5-trimethylhydroquinone and the ester formed as the further product. 2,3,5-Trimethylhydroquinone can be converted into (all-rac)- α -tocopherol by known procedures by reaction with isophytol, preferably in a biphasic solvent system, e.g. in a solvent system comprising a polar solvent such as ethylene or propylene carbonate, and a non-polar solvent, particularly an aliphatic hydrocarbon such as heptane, see, e.g. international application PCT/EP03/01556.

The invention is illustrated further by the following Examples.

15 Example 1

A 50-ml four-necked flat-bottomed flask equipped with a thermometer, a glass-tube (\varnothing 5 mm) for Ar-purge, a reflux condenser and a magnetic stirring bar was charged with methanetrissulfonic acid (see Table 1 below) and 10.324 g (66 mmol) of ketoisophorone. Within 2 min, acetic anhydride was added dropwise (see Table 1 below) under rapid stirring. During addition, the mixture turned dark yellow to finally dark brown and the internal temperature increased. After cooling to the desired reaction temperature that temperature was maintained by means of an oil bath. Samples were withdrawn and submitted to qualitative GC-analysis. After the reaction time (see Table 1 below), the reaction mixture was cooled to room temperature and the catalyst was deactivated by addition of 3.7 g (70 mmol) anhydrous sodium carbonate. The reaction mixture was concentrated by 40°C/10mbar thereby distilling off acetic acid and unreacted anhydride. The crude product was analyzed by GC using squalane as internal standard. The results and reactions conditions are given in Table 1 below:

Table 1

(Ac ₂ O) [mmol]	(SO ₃ H) ₃ CH [mg]	(SO ₃ H) ₃ CH [mol %]	T [°C]	time [h]	conversion [%]	TMHQ-DA [%]
200	171.2	1.0	25	4	97.2	91.8
333	171.2	1.0	25	4	99.3	91.1
666	171.2	1.0	25	4	99.2	94.0
200	93.5	0.55	25	22	78.6	69.0
200	171.2	1.0	25	22	100	96.6
200	342.4	2.0	25	22	100	93.2
200	120	0.7	60	4	99.1	91.5
200	120	0.7	60	4	99.3	92.9
200	120	0.7	60	4	98.9	92.3
200	34.2	0.2	70	4	83.5	72.2
200	94.2	0.55	70	4	100	92.6
200	154.1	0.9	70	4	100	94.2
200	171.2	1.0	40	4.5	98.8	88.2
200	342.4	2.0	40	3.5	100	90.0

5 The values given are averages from several measurements (two or three) and two experiments.

Ac₂O : acetic anhydride; TMHQ-DA : 2,3,5-trimethyl-1,4-hydroquinone diacetate [yield]

Example 2

- 10 Using a 230 ml-flask the procedure of Example 1 was repeated with changing ratios of ketoisophorone and acetic anhydride. The reaction conditions and results are given in Table 2 below:

Table 2

(Ac ₂ O) [mmol]	KIP/Ac ₂ O [ratio]	T [°C]	time [h]	conversion [%]	TMHQ-DA [%]	S(TM HQ-DA) [%]
299	1 : 2.25	45	12	89.7	85.8	95.6
332	1 : 2.5	45	12	91.9	87.2	94.9
398	1 : 3	45	12	95.6	90.9	95.1
663	1 : 5	45	12	98.6	94.5	95.8
1327	1 : 10	45	12	98.1	93.0	94.8
398	1 : 3	25	24	83.6	80.6	96.3

- 15 The values given are averages from several measurements (two or three) and two experiments. S : Selectivity.

Example 3

In analogy to the procedure of Example 1, the reaction conditions were optimized using a statistical model (STAVEX). 20.32 g (200 mmol) of acetic anhydride were added within 10 min. to 10.324 g (66 mmol) of ketoisophorone. The results are tabulated in Table 3 below:

Table 3

(SO ₃ H) ₃ CH [mg]	(SO ₃ H) ₃ CH [mmol]	(SO ₃ H) ₃ CH [mol %]	T [°C]	time [h]	conversion [%]	TMHQ-DA [%]
17.2	0.07	0.1	25	4	0	0
17.4	0.07	0.1	100	4	43.2	14.8
172.5	0.67	1.0	100	4	100	83.5
85.6	0.33	0.5	40	14	97.6	89.7
17.2	0.07	0.1	25	24	10.8	1.0
85.5	0.33	0.5	55	14	100	90.3
85.5	0.33	0.5	55	22	100	91.4
171.4	0.67	1.0	25	4	83.9	75.6
85.6	0.33	0.5	85	14	100	89.6
17.1	0.07	0.1	100	24	45.6	16.4
34.4	0.13	0.2	55	14	88.7	79.0
85.5	0.33	0.5	55	6	99.5	90.4
171.1	0.67	1.0	25	24	99.3	92.7
154.2	0.60	0.9	55	14	100	91.3
171.4	0.67	1.0	100	24	100	84.6
171.3	0.67	1.0	60	7	100	91.2
128.1	0.50	0.8	60	10	100	91.9
85.5	0.33	0.5	60	12	100	93.5

10 Example 4

Based on the results of Example 3, experiments under optimized conditions were carried out. The results are summarized in Table 4

Table 4

(Ac ₂ O) [mmol]	KIP/ Ac ₂ O ratio	(SO ₃ H) ₃ CH [mol %]	T [°C]	time [h]	conversion [%]	TMHQ-DA [%]	S(TMHQ-DA) [%]
398	1 : 3	0.5	60	6	97.4	92.0	94.5
663	1 : 5	1.0	25	24	100	96.7	96.7

What is claimed is:

1. Process for the preparation of 2,3,5-trimethyl-1,4-hydroquinone diacylates by reacting 3,5,5-trimethyl-1,4-benzoquinone with an acylating agent in the presence of methane
5 trisulfonic acid.
2. A process as in claim 1 wherein the reaction is carried out in the presence of about 0.01 mol% to about 2 mol% of methane trisulfonic acid, based on 3,5,5-trimethyl-1,4-benzoquinone.
10
3. A process as in claim 2 wherein the reaction is carried out in the presence of about 0.1 mol% to about 1 mol% of methane trisulfonic acid, based on 3,5,5-trimethyl-1,4-benzoquinone.
- 15 4. A process as in any one of claims 1-3 wherein the molar ratio of acylating agent to 3,5,5-trimethyl-1,4-benzoquinone to is about 10 to 1, preferably about 3 to 1.
5. A process as in any one of claims 1-4 wherein the acylating agent is acetic anhydride.
- 20 6. A process as in any one of claims 1-5 wherein the reaction is carried out at about 0 °C to about 140 °C, preferably at about 20 °C to about 70 °C.
7. A process as in any one of claims 1-6 wherein the 2,3,5-trimethyl-1,4-hydroquinone diester obtained is converted in a manner known per se to α -tocopherol.

INTERNATIONAL SEARCH REPORT

International Application No
EP2004/012058

A. CLASSIFICATION OF SUBJECT MATTER

IPC 7 C07C67/08 C07C69/017 C07C69/16 C07D311/72

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 C07C C07D

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the International search (name of data base and, where practical, search terms used)

EPO-Internal, CHEM ABS Data

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	DE 21 49 159 A (EASTMAN KODAK CO.) 6 April 1972 (1972-04-06) cited in the application claims 1-9	1-7
Y	US 6 103 924 A (N. SHI ET AL.) 15 August 2000 (2000-08-15) claims 1-21	1-7
Y	WO 03/051812 A (ROCHE VITAMINS AG) 26 June 2003 (2003-06-26) cited in the application claims 1-14	1-7
Y	US 3 053 884 A (G. P. TOUEY, R. H. GOINS) 11 September 1962 (1962-09-11) claims 1-8	1-7
	-/-	

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

* Special categories of cited documents :

A document defining the general state of the art which is not considered to be of particular relevance

E earlier document but published on or after the international filing date

L document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

O document referring to an oral disclosure, use, exhibition or other means

P document published prior to the international filing date but later than the priority date claimed

T later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

X document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

Y document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

* & * document member of the same patent family

Date of the actual completion of the international search

14 February 2005

Date of mailing of the international search report

25/02/2005

Name and mailing address of the ISA

European Patent Office, P.B. 5618 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

Herz, C

INTERNATIONAL SEARCH REPORT

International Application No

P/EP2004/012058

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	M. SCHNEIDER ET AL.: "Industrial application of Nafion-systems in rearrangement-aromatisation, transesterification, alkylation, and ring-closure reactions" APPLIED CATALYSIS A GENERAL, vol. 220, no. 1-2, 2001, pages 51-58, XP002317570 the whole document	1-7
Y	----- US 2002/004619 A1 (S. KRILL, K. HUTHMACHER) 10 January 2002 (2002-01-10) claims 1-14	1-7
Y	----- EP 0 850 910 A (DAICEL CHEMICAL INDUSTRIES, LTD.) 1 July 1998 (1998-07-01) claims 1-8	1-7
Y	----- DE 198 05 690 A (DEGUSSA-HÜLS) 19 August 1999 (1999-08-19) claims 1-11 -----	1-7

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

EP2004/012058

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
DE 2149159	A	06-04-1972	DE 2149159 A1	06-04-1972
			US T900015 I4	25-07-1972
US 6103924	A	15-08-2000	DE 19817644 A1	28-10-1999
			BR 9902048 A	02-05-2000
			CA 2269404 A1	21-10-1999
			CN 1234389 A	10-11-1999
			DE 59909987 D1	26-08-2004
			DK 952137 T3	08-11-2004
			EP 0952137 A1	27-10-1999
			ID 22456 A	21-10-1999
			IL 129529 A	25-11-2001
			IN 185858 A1	12-05-2001
			JP 11349533 A	21-12-1999
			TW 490459 B	11-06-2002
WO 03051812	A	26-06-2003	AU 2002356637 A1	30-06-2003
			WO 03051812 A1	26-06-2003
			EP 1463708 A1	06-10-2004
US 3053884	A	11-09-1962	NONE	
US 2002004619	A1	10-01-2002	DE 10017494 A1	11-10-2001
			AT 259772 T	15-03-2004
			AU 4835701 A	23-10-2001
			CA 2399248 A1	18-10-2001
			CN 1420859 T	28-05-2003
			DE 60102078 D1	25-03-2004
			DE 60102078 T2	23-12-2004
			DK 1268390 T3	10-05-2004
			WO 0177058 A1	18-10-2001
			EP 1268390 A1	02-01-2003
			JP 2003530377 T	14-10-2003
			MX PA02009115 A	12-03-2003
EP 0850910	A	01-07-1998	CN 1186065 A ,C	01-07-1998
			DE 69708450 D1	03-01-2002
			DE 69708450 T2	16-05-2002
			EP 0850910 A1	01-07-1998
			JP 10237020 A	08-09-1998
			US 5908956 A	01-06-1999
DE 19805690	A	19-08-1999	DE 19805690 A1	19-08-1999
			BR 9900655 A	09-05-2000
			CA 2261404 A1	12-08-1999
			EP 1008584 A1	14-06-2000
			ID 21942 A	12-08-1999
			IN 186229 A1	14-07-2001
			JP 11292812 A	26-10-1999
			US 5969176 A	19-10-1999